

Possible Association of the Ubiquitin-Specific Peptidase 46 Gene (*USP46*) with Affective Temperamental Traits in Healthy Korean Volunteers

Young Jun Boo^{1,2}, Chun Il Park^{1,2}, Hae Won Kim^{2,3}, Se Joo Kim^{1,2} ✉, and Jee In Kang^{1,2} ✉

¹Department of Psychiatry, Yonsei University College of Medicine, Seoul, Republic of Korea

²Institute of Behavioral Science in Medicine, Yonsei University College of Medicine, Seoul, Republic of Korea

³Department of Medical Education, Yonsei University College of Medicine, Seoul, Republic of Korea

Objective Ubiquitin-specific peptidase 46 gene (*USP46*) polymorphisms is part of ubiquitin-proteasome system, which is responsible for dynamic cellular processes such as the regulation of cell cycle. *USP46* has been reported to be associated with major depressive disorder. The objective of the present study was to investigate the association of *USP46* polymorphisms with affective temperamental traits in healthy subjects.

Methods A total of 557 Korean healthy volunteers were recruited, and 545 subjects (328 male, 217 female) were included in the final analysis. The DNA of the subjects was isolated from saliva samples. Two single-nucleotide polymorphisms (SNPs) rs346005, rs2244291 in *USP46* were genotyped. Affective temperaments were assessed using the Korean version of Temperament Evaluation of the Memphis, Pisa, Paris, and San Diego Autoquestionnaire (TEMPS-A).

Results A significant association was found between rs346005 genotypes and TEMPS-A only in male subjects. In particular, subjects with the CC genotype of rs346005 showed a more depressive temperament than subjects with AA or CA genotypes in males. For rs2244291, there were no associations between the rs2244291 genotypes and TEMPS-A scores.

Conclusion Some affective temperaments may serve as a genetic predisposing factors for affective disorders, such as depressive disorder, via vulnerability genes related to the ubiquitin-proteasome system.

Psychiatry Investig 2019;16(1):87-92

Key Words Affective temperament, Genetic association study, TEMPS-A, Ubiquitin-proteasome system, *USP46*.

INTRODUCTION

Affective temperaments refer to characteristic patterns in the quality and intensity of baseline mood levels and emotional and behavioral responses and regulations to internal and external stimuli, which tend to appear early in life and be

temporally stable.¹⁻³ Akiskal et al.¹ proposed five affective temperamental traits: depressive, cyclothymic, hyperthymic, irritable, and anxious temperaments and developed a valid assessment scale for affective temperaments, viz. the Temperament Evaluation of the Memphis, Pisa, Paris, and San Diego Autoquestionnaire (TEMPS-A). These affective temperaments are not only considered to be subclinical manifestation of affective disorders,^{1,4,5} but are also known to play a crucial role in the clinical prognosis of psychiatric illnesses.⁶⁻⁹ A recent study of affective temperament and suicide showed that an anxious temperament is a strong risk factor for suicidal attempts, regardless of the presence of mental disorders, while depressive and irritable temperaments are risk factors for suicidal attempts in patients with mental disorders.⁶ Therefore, a better understanding of affective temperaments is crucial to the understanding of the development of and treatments for mental disorders.

Affective temperaments are known to be moderately heri-

Received: April 30, 2018 Revised: August 14, 2018

Accepted: October 2, 2018

✉ Correspondence: Se Joo Kim, MD, PhD

Department of Psychiatry and Institute of Behavioral Science in Medicine, Yonsei University College of Medicine, 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, Republic of Korea

Tel: +82-2-2228-1620, Fax: +82-2-313-0890, E-mail: kimsejoo@yuhs.ac

✉ Correspondence: Jee In Kang, MD, PhD

Department of Psychiatry and Institute of Behavioral Science in Medicine, Yonsei University College of Medicine, 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, Republic of Korea

Tel: +82-2-2228-1620, Fax: +82-2-313-0890, E-mail: jeeinkang@yuhs.ac

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

table, with heritability ranging from 21% to 52% for the five affective temperaments in 101 bipolar families.¹⁰ Additionally, several candidate gene association studies have been conducted to identify genes for affective temperaments. Previous genetic association approaches of affective temperaments usually targeted monoaminergic systems, including serotonin-related genes,^{11–14} and dopaminergic genes.¹³ A genome-wide association study has revealed several variants that are significantly associated with a specific affective temperament in bipolar patients.¹⁵ Since affective temperaments may be quantitative endophenotypes for affective disorders, representing genetic heterogeneity, the interest in genetic studies of affective temperaments has increased.

Besides the monoaminergic genes, the ubiquitin-proteasome system (UPS), a major protein degradation pathway, which is responsible for dynamic cellular processes, such as the regulation of cell cycle and the response to genotoxic and proteotoxic stress,¹⁶ may be a potential candidate system contributing to affective temperaments as well as affective disorders. The UPS is a critical contributor to the construction of brain networks via extensive regulation of synaptic plasticity.^{17,18} In recent years, synaptic plasticity has emerged as a component of the pathophysiology of depression.^{19,20} Interestingly, some studies have revealed direct associations between the UPS and depressive disorders.^{21–23} In an animal study using quantitative trait locus mapping, the ubiquitin-specific peptidase 46 gene (*USP46*) was proposed to be responsible for depressive-like behavior in mice in depression-inducing tests, such as tail suspension and forced swimming tests.²³ Furthermore, *USP46* knock-out mice also showed differences in depression-inducing test.²² Moreover, a human study showed significant differences in *USP46* single-nucleotide polymorphisms (SNPs) between patients with depressive disorder and normal controls.²¹ Taken together, these findings may indicate that *USP46* may be a candidate gene involved in the development of affective temperaments, particularly in depressive temperaments, as a subclinical manifestation of affective disorders.

The objective of the present study was to investigate the association of *USP46* polymorphisms with affective temperamental traits in healthy subjects.

METHODS

Subjects

A total of 557 healthy Korean volunteers (334 males, 223 females) were recruited using advertisements. Most of the subjects were college students. All participants were interviewed by psychiatrists, and subjects with any current or previous Axis I psychiatric disorders according to the Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (DSM-IV), a

family history of psychiatric disorders, or neurologic disorders were excluded. Twelve subjects (2.16% of the total subjects) were excluded based on the exclusion criteria. A total of 545 subjects (328 male, 217 female) were included in the final analysis. All subjects provided informed consent before the study. The study protocol was approved by the Institutional Review Board of Severance Hospital (Severance IRB 4-2010-0577).

SNP selection

To date, only one study has reported positive association between *USP46* gene and psychiatric phenotypes. Fukuo et al.²¹ found significant association between haplotypes consisting of three (rs2244291–rs10034164–rs346005) or four (rs12646800–rs2244291–rs10034164–rs346005) SNPs of seven tagging SNPs ($r^2 \geq 0.8$) capturing 30 SNPs (minor allele frequency; $MAF \geq 5$) covering the *USP46* gene region.²¹ Therefore, we selected SNPs of rs12646800, rs2244291, rs10034164, and rs346005 in this study. However, we further excluded rs12646800 and rs10034164, as $MAFs$ were less than 0.1 (0.056 for rs12646800, 0.087 for rs10034164). Especially, there were no minor homozygotes for rs10034164 in the HapMap database (release #27, population: Japanese in Tokyo). Finally, we included two SNPs (rs2244291 and rs346005) in this study. SNP rs2244291 is located in intron, and SNP rs346005 is located 2 kb upstream of *USP46* start site (*USP46*; 52590960–52659335, rs346005; 52588572).

Genotyping

The DNA of the subjects was isolated from saliva collected in OG-500 Oragene® self-collection kit (tube format “OG-500”), using standard techniques. Genotyping of *USP46* SNPs was performed using a single-base primer extension assay (ABI PRISM® SNaPshot™ Multiplex kit; ABI, Foster City, CA, USA) according to the manufacturer's instructions. Genotyping was performed by DNA Link, Inc. (Seoul, South Korea), in a manner blinded to any other study data.

Assessments

Affective temperament was assessed with the Korean version of the Memphis, Pisa, Paris, and San Diego Autoquestionnaire (TEMPS-A, a self-reported temperament scale for evaluating temperamental dimensions, with five subscales, including depressive, cyclothymic, hyperthymic, irritable, and anxious temperaments.¹ This scale consists of 110 “yes-or-no” type questions in total, of which 109 are applicable to male subjects. All participants completed the questionnaire.

Statistical analyses

Previous combined analysis showed the differences in TEMPS-A subscales scores between males and females.²⁴ Therefore, all the analyses in this study were conducted separately by

gender. Multiple analyses of covariance (MANCOVA) were performed using the total scores for each TEMPS-A affective temperament dimension as dependent variables, and *USP46* SNP genotypes as fixed factors. Age was controlled as a covariate to control possible effects. The two SNPs were analyzed separately for each gender. Further analyses were conducted for significant associations, with Fisher's least significant difference post-hoc tests to evaluate differences among genotypes. Data analysis was performed using SPSS 23.0 for Windows (IBM Corp., Armonk, NY, USA). The statistical significance was set at $p < 0.05$ for all tests.

Table 1. Demographic characteristics of the subjects

	Male (N=328)	Female (N=217)	p
Age (years)	22.55 (2.64)	21.57 (2.47)	<0.001
Education (years)	14.32 (1.46)	14.39 (1.58)	0.596
TEMPS-A			
Depressive	6.73 (3.03)	7.51 (2.90)	0.003
Cyclothymic	6.83 (4.56)	8.03 (4.57)	0.003
Hyperthymic	11.54 (4.79)	10.79 (4.54)	0.068
Irritable	3.60 (3.27)	3.73 (3.45)	0.633
Anxious	5.56 (5.19)	7.28 (5.42)	<0.001

Mean (SD). SD: standard deviation, TEMPS-A: Temperament Evaluation of the Memphis, Pisa, Paris, and San Diego Autoquestionnaire

Table 2. Multivariate analysis comparing scores for each affective temperament dimension for *USP46* single-nucleotide polymorphism rs346005 (mean±SEM) after controlling for age

	Male (N=328)			P	Female (N=217)			P
	AA (N=97)	CA (N=161)	CC (N=70)		AA (N=65)	CA (N=111)	CC (N=41)	
MANCOVA	Wilks $\lambda=0.929$, $F_{(10,640)}=2.384$, $p=0.009$				Wilks $\lambda=0.979$, $F_{(10,418)}=0.450$, $p=0.921$			
Depressive	6.79±0.30	6.20±0.23	7.87±0.36	<0.001	7.68±0.36	7.28±0.28	7.86±0.45	0.472
Cyclothymic	6.47±0.46	6.78±0.36	7.45±0.54	0.380	8.34±0.58	7.96±0.44	7.71±0.72	0.886
Hyperthymic	11.46±0.48	11.91±0.38	10.80±0.57	0.258	10.55±0.57	10.93±0.44	10.78±0.71	0.869
Irritable	3.34±0.33	3.43±0.26	4.34±0.39	0.097	3.59±0.44	3.69±0.33	4.10±0.54	0.750
Anxious	5.97±0.53	4.97±0.41	6.36±0.62	0.120	7.53±0.68	7.16±0.52	7.22±0.85	0.906

SEM: standard error of mean, MANCOVA: multiple analyses of covariance

Table 3. Multivariate analysis comparing scores for each affective temperament dimension for *USP46* gene single-nucleotide polymorphism rs2244291 (mean±SEM) after controlling for age

	Male (N=328)			P	Female (N=217)			P
	AA (N=198)	GA (N=118)	GG (N=12)		AA (N=138)	GA (N=68)	GG (N=11)	
MANCOVA	Wilks $\lambda=0.966$, $F_{(10,640)}=1.124$, $p=0.341$				Wilks $\lambda=0.935$, $F_{(10,418)}=1.421$, $p=0.168$			
Depressive	6.62±0.22	6.94±0.28	6.44±0.88	0.643	7.56±0.25	7.27±0.35	8.32±0.88	0.505
Cyclothymic	6.62±0.32	7.37±0.42	5.06±1.31	0.146	8.17±0.39	7.61±0.56	8.80±1.38	0.601
Hyperthymic	11.64±0.34	11.36±0.44	11.56±1.38	0.879	10.28±0.39	11.84±0.55	10.64±1.36	0.070
Irritable	3.38±0.23	3.99±0.30	3.33±0.95	0.266	3.57±0.29	3.82±0.42	5.27±1.04	0.287
Anxious	5.63±0.37	5.67±0.48	3.39±1.50	0.339	6.92±0.46	7.61±0.66	9.84±1.63	0.192

SEM: standard error of mean, MANCOVA: multiple analyses of covariance

RESULTS

Baseline characteristics

Baseline demographic data and characteristics, including TEMPS-A scores, are summarized in Table 1. The mean age of the participants was 22.55 years (standard deviation; SD=2.64 years) for males and 21.57 years (SD=2.47 years) for females. The mean education level of the participants were 14.32 years (SD=1.46 years) for males and 14.39 years (SD=1.58 years) for females. There were significant differences in age ($p < 0.001$) and depressive ($p = 0.003$), cyclothymic ($p = 0.003$), and anxious ($p < 0.001$) dimension scores between male subjects and female subjects.

Association analyses between *USP46* SNPs and five affective temperamental dimensions

The genotype distribution of the two *USP46* SNPs, rs346005 and rs2244291 were in accordance with the Hardy-Weinberg equilibrium ($\chi^2 = 0.068$, $p = 0.794$ for rs346005, $\chi^2 = 0.161$, $p = 0.6835$ for rs2244291).

MANCOVA results are presented in Table 2 through Table 3. The MANCOVA showed a significant association between TEMPS-A and rs346005 genotypes in male subjects (Wilks $\lambda = 0.929$, $F_{(10,640)} = 2.384$, $p = 0.009$). Among temperament dimensions, there was a significant difference in depressive temperament among the genotypes of rs346005 in male sub-




Figure 1. Post-hoc analysis comparing the TEMPS-A depressive subscale score for *USP46* single-nucleotide polymorphism rs346005. Error bars indicate the standard error of mean. TEMPS-A: Temperament Evaluation of the Memphis, Pisa, Paris, and San Diego Autoquestionnaire.

jects ($F=7.79$, $p<0.001$, $\eta^2=0.046$). The association was not observed in female subjects ($F=0.75$, $p=0.472$, $\eta^2=0.007$) (Table 2). In contrast, for rs2244291, no associations were found between the genotypes and the five affective temperaments, in males or females (Table 3).

Post-hoc comparisons showed that subjects who had the CC genotype (mean=7.89, SD=0.45) at rs346005 showed a higher depressive temperament score than that of subjects with the AA genotype (mean=6.80, SD=0.30, $p<0.05$) and the CA genotype (mean=6.18, SD=0.21, $p<0.0001$) in male subjects. There is no significant difference in the depressive temperament score between subjects with the AA genotype and subjects with the CA genotype ($p=0.103$) (Figure 1).

DISCUSSION

We investigated the associations of the *USP46* polymorphisms with the affective temperamental traits in healthy college students. The major finding was that the CC genotype of *USP46* rs346005 was related to a higher depressive temperament score in male subjects. These findings suggest that affective temperamental traits are influenced by *USP46*.

Genetic associations between the ubiquitin proteasome system and affective temperament have not yet been reported. Although there is little direct evidence, our finding of the involvement of *USP46* in depressive temperament is supported by indirect evidence. Theoretically, deubiquitinating enzymes, such as *USP46*, could play a role in regulating the stability of cellular protein and synaptic trafficking.²⁵ Animal studies have shown that *USP46*-mutated mice exhibit a significantly

shorter immobility time than wild type mice in a tail suspension test (TST), which indicates a change in depressive state.²³ Furthermore, a genetic association study of major depression showed significant involvement of a *USP46* haplotype in a Japanese population with major depressive disorder.²¹ In the study, rs346005 was the only SNP that showed significant association with major depressive disorder. Considering that a depressive temperament is a subclinical form of major depressive disorder, it is possible that *USP46* may be involved in the pathophysiology of depression.

The biological mechanism underlying the association between *USP46* and depressive temperament remains unclear. However, some possible explanations of the association can be proposed. First, *USP46* may act as a mood regulator by modulating neurotransmitter systems, such as serotonin and GABA. The antidepressant imipramine, a tricyclic antidepressant that inhibits reuptake of monoamines, such as serotonin and norepinephrine, reduced immobility in the TST of *USP46* mutated mice,²³ which supports the monoamine hypothesis.²⁶ In contrast, another study using *USP46*-knock-out mice reported that depressive behaviors in these mice were changed by administration of nitrazepam, GABA_A receptor agonist,²² supporting the GABA hypothesis of major depressive disorder.²⁷ Second, the ubiquitin system is known to regulate the circadian system by controlling circadian clock protein.²⁸ Associations between the circadian system and affective temperament have previously been reported.^{29,30} In a previous study, a depressive temperamental trait appeared to be associated with evening-type circadian preferences, along with cyclothymic, irritable, and anxious temperaments.³⁰ Therefore, it can be assumed that *USP46* influences affective temperamental traits by modulating the circadian system. Third, *USP46* may affect the affective temperament by modulating synaptic plasticity. *USP46* has been reported to play a role in regulating alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid receptor (AMPA), one of the key receptors involved in long-term potentiation and depression.²⁵ *USP46*, which is known to be enriched at the neural synapse, may modulate brain function by deubiquitinating AMPAR. Because evidence supporting these speculations is lacking, further studies elucidating the pathways that may mediate the association between UPS and depressive temperament should be performed.

Unlike in male subjects, no main effects of *USP46* polymorphisms were observed in female subjects. This finding could be explained in several ways. In general, gene expression and splicing is known to differ between males and females.³¹ From other perspectives, this gender difference might be related to an effect of sex hormones. Several studies have reported that the sex hormone estrogen modulates the ubiquitin systems,^{32,33} and plays an important role in synaptic plasticity.³⁴

In this study, there was no difference in genotype distribution of rs346005 between males and females. This suggested that *USP46* rs346005 is not the major cause of difference in depressive temperament between males and females. In addition, considering the fact that complex traits, such as affective temperaments, are mediated by many genes (not one gene) and their interactions, the higher score of depressive temperament in females seems to be affected by other conditions, such as sexual hormone and/or X-chromosome. We could not find other studies showing the sexual dimorphisms of *USP46* rs346005, since there is only a few existing studies on *USP46*. However, we found several other evidence, although not *USP46*, suggesting genetic sexual dimorphisms on temperamental traits. Stankova et al.³⁵ reported that the sexual dimorphic effects of oxytocin receptor gene (*OXTR*) showed gender difference on harm avoidance; one of the aspects of temperamental traits. In our previous study of dopamine receptor D4 (*DRD4*) gene on affective temperament, the association between *DRD4* polymorphism and affective temperament (cyclothymic and irritable) was found only in male subjects.¹³ In addition, since this study involved a small number of female subjects (n=217), the small sample size may not have been sufficient to obtain significant results.

There are several limitations in this study. First, most of our data were collected from college students, and it may not be generalizable. Although our sample can be considered a homogeneous group, with similar age, educational levels, and ethnicity, the results of this study should be considered cautiously. The results need to be replicated in other populations of different ages and ethnicities. Second, the present study was performed in a relatively small number of subjects. A further study in a larger sample size is needed to confirm the results. Third, we assessed only two SNPs. There are many more SNPs in *USP46*, and thus a more precise result could be obtained if additional SNPs were to be assessed. Fourth, affective temperamental scores could have been biased by social desirability, since TEMPS-A is a self-reporting scale. Fifth, we did not analyze gene-gene interaction and gene-environment interaction. Despite these limitations, this is the first study that reveals a possible genetic association between the UPS and affective temperaments.

In summary, the present study showed an association between a SNP in *USP46* and depressive temperament. Our findings suggest that some affective temperaments may be genetically predisposed to developing into affective disorders via vulnerable genes related to the UPS. In particular, *USP46* may play a role in the pathophysiology of major depressive disorders by modulating a depressive temperament through dynamic proteolysis processes involving the UPS. Further research is required to elucidate influence of *USP46*

polymorphisms of affective temperaments in patients with mood disorders.

Acknowledgments

This study was supported by a grant from the Disease Oriented Translational Research through the Korea Health Industry Development Institute (KHIDI), South Korea (grant number: HI14C0202).

REFERENCES

1. Akiskal HS, Akiskal KK, Haykal RF, Manning JS, Connor PD. TEMPS-A: progress towards validation of a self-rated clinical version of the Temperament Evaluation of the Memphis, Pisa, Paris, and San Diego Autoquestionnaire. *J Affect Disord* 2005;85:3-16.
2. Kawamura Y, Akiyama T, Shimada T, Minato T, Umekage T, Noda Y, et al. Six-year stability of affective temperaments as measured by TEMPS-A. *Psychopathology* 2010;43:240-247.
3. Nigg JT. Temperament and developmental psychopathology. *J Child Psychol Psychiatry* 2006;47:395-422.
4. Kraepelin E. Manic-Depressive Insanity and Paranoia. Edinburgh: E. & S. Livingstone; 1921.
5. Toda H, Inoue T, Tsunoda T, Nakai Y, Tanichi M, Tanaka T, et al. Affective temperaments play an important role in the relationship between childhood abuse and depressive symptoms in major depressive disorder. *Psychiatry Res* 2016;236:142-147.
6. Karam EG, Itani L, Fayyad J, Hantouche E, Karam A, Mneimneh Z, et al. Temperament and suicide: a national study. *J Affect Disord* 2015;184:123-128.
7. Pompili M, Innamorati M, Gonda X, Serafini G, Sarno S, Erbutto D, et al. Affective temperaments and hopelessness as predictors of health and social functioning in mood disorder patients: a prospective follow-up study. *J Affect Disord* 2013;150:216-222.
8. Takeshima M, Oka T. Comparative analysis of affective temperament in patients with difficult-to-treat and easy-to-treat major depression and bipolar disorder: possible application in clinical settings. *Compr Psychiatry* 2016;66:71-78.
9. Vazquez GH, Gonda X, Zaratiegui R, Lorenzo LS, Akiskal K, Akiskal HS. Hyperthymic temperament may protect against suicidal ideation. *J Affect Disord* 2010;127:38-42.
10. Greenwood TA, Badner JA, Byerley W, Keck PE, McElroy SL, Remick RA, et al. Heritability and genome-wide SNP linkage analysis of temperament in bipolar disorder. *J Affect Disord* 2013;150:1031-1040.
11. Borkowska A, Bielinski M, Szczesny W, Szwed K, Tomaszewska M, Kalwa A, et al. Effect of the 5-HTTLPR polymorphism on affective temperament, depression and body mass index in obesity. *J Affect Disord* 2015;184:193-197.
12. Gonda X, Rihmer Z, Zsombok T, Bagdy G, Akiskal KK, Akiskal HS. The 5HTTLPR polymorphism of the serotonin transporter gene is associated with affective temperaments as measured by TEMPS-A. *J Affect Disord* 2006;91:125-131.
13. Kang JI, Namkoong K, Kim SJ. The association of 5-HTTLPR and *DRD4* VNTR polymorphisms with affective temperamental traits in healthy volunteers. *J Affect Disord* 2008;109:157-163.
14. Landaas ET, Johansson S, Halmoy A, Oedegaard KJ, Fasmer OB, Haavik J. No association between the serotonin transporter gene polymorphism 5-HTTLPR and cyclothymic temperament as measured by TEMPS-A. *J Affect Disord* 2011;129:308-312.
15. Greenwood TA, Akiskal HS, Akiskal KK; Bipolar Genome S, Kelsoe JR. Genome-wide association study of temperament in bipolar disorder reveals significant associations with three novel Loci. *Biol Psychiatry* 2012;72:303-310.
16. Craney A, Rape M. Dynamic regulation of ubiquitin-dependent cell cycle control. *Curr Opin Cell Biol* 2013;25:704-710.
17. Hamilton AM, Zito K. Breaking it down: the ubiquitin proteasome sys-

- tem in neuronal morphogenesis. *Neural Plast* 2013;2013:196848.
18. Hegde AN, Haynes KA, Bach SV, Beckelman BC. Local ubiquitin-proteasome-mediated proteolysis and long-term synaptic plasticity. *Front Mol Neurosci* 2014;7:96.
19. Duman RS, Aghajanian GK, Sanacora G, Krystal JH. Synaptic plasticity and depression: new insights from stress and rapid-acting antidepressants. *Nat Med* 2016;22:238-249.
20. Wainwright SR, Galea LA. The neural plasticity theory of depression: assessing the roles of adult neurogenesis and PSA-NCAM within the hippocampus. *Neural Plast* 2013;2013:805497.
21. Fukuo Y, Kishi T, Kushima I, Yoshimura R, Okochi T, Kitajima T, et al. Possible association between ubiquitin-specific peptidase 46 gene and major depressive disorders in the Japanese population. *J Affect Disord* 2011;133:150-157.
22. Imai S, Mamiya T, Tsukada A, Sakai Y, Mouri A, Nabeshima T, et al. Ubiquitin-specific peptidase 46 (*Usp46*) regulates mouse immobile behavior in the tail suspension test through the GABAergic system. *PLoS One* 2012;7:e39084.
23. Tomida S, Mamiya T, Sakamaki H, Miura M, Aosaki T, Masuda M, et al. *Usp46* is a quantitative trait gene regulating mouse immobile behavior in the tail suspension and forced swimming tests. *Nat Genet* 2009;41:688-695.
24. Vazquez GH, Tondo L, Mazzarini L, Gonda X. Affective temperaments in general population: a review and combined analysis from national studies. *J Affect Disord* 2012;139:18-22.
25. Huo Y, Khatri N, Hou Q, Gilbert J, Wang G, Man HY. The deubiquitinating enzyme *USP46* regulates AMPA receptor ubiquitination and trafficking. *J Neurochem* 2015;134:1067-1080.
26. Hirschfeld RM. History and evolution of the monoamine hypothesis of depression. *J Clin Psychiatry* 2000;61(Suppl 6):4-6.
27. Luscher B, Shen Q, Sahir N. The GABAergic deficit hypothesis of major depressive disorder. *Mol Psychiatry* 2011;16:383-406.
28. Stojkovic K, Wing SS, Cermakian N. A central role for ubiquitination within a circadian clock protein modification code. *Front Mol Neurosci* 2014;7:69.
29. Chrobak AA, Tereszko A, Dembinska-Krajewska D, Arciszewska A, Siwek M, Dudek D, et al. Morningness-eveningness and affective temperaments assessed by the Temperament Evaluation of Memphis, Pisa and San Diego-Autoquestionnaire (TEMPS-A). *Chronobiol Int* 2017;34:57-65.
30. Park CI, An SK, Kim HW, Koh MJ, Namkoong K, Kang JJ, et al. Relationships between chronotypes and affective temperaments in healthy young adults. *J Affect Disord* 2015;175:256-259.
31. Trabzuni D, Ramasamy A, Imran S, Walker R, Smith C, Weale ME, et al. Widespread sex differences in gene expression and splicing in the adult human brain. *Nat Commun* 2013;4:2771.
32. Lai YJ, Liu L, Hu XT, He L, Chen GJ. Estrogen Modulates *ubc9* Expression and Synaptic Redistribution in the Brain of APP/PS1 Mice and Cortical Neurons. *J Mol Neurosci* 2017;61:436-448.
33. Ogawa M, Yamaji R, Higashimura Y, Harada N, Ashida H, Nakano Y, et al. 17 β -estradiol represses myogenic differentiation by increasing ubiquitin-specific peptidase 19 through estrogen receptor α . *J Biol Chem* 2011;286:41455-41465.
34. Spencer-Segal JL, Tsuda MC, Mattei L, Waters EM, Romeo RD, Milner TA, et al. Estradiol acts via estrogen receptors α and β on pathways important for synaptic plasticity in the mouse hippocampal formation. *Neuroscience* 2012;202:131-146.
35. Stankova T, Eichhammer P, Langguth B, Sand PG. Sexually dimorphic effects of oxytocin receptor gene (*OXTR*) variants on Harm Avoidance. *Biol Sex Differ* 2012;3:17.